

Soil Feeding Schedule

SOIL & SOILLESS FEEDING TIPS: Use FoxFarm Ocean Forest® Potting Soil when transplanting seedlings into larger containers. For best results feed with every other watering. During the flowering stage reduce the amount of light from 18 hours to 12 hours. Maintain a pH of 6.3 to 6.8 to prevent nutrient lock up and reduce stress on plants. Use primary nutrients for abundant growth and stocky, robust plants. For high-octane yields add supplements to the weekly diet. Never mix pure concentrates together, always add water first.

Should plants show signs of stress or color irregularities, flush your system with FoxFarm BushDoctor® SledgeHammer® combined with FoxFarm Big Bloom®. Use twice the volume of water as a typical feeding to remove unwanted salt build-up and heal the root bio-culture. For edible plants flush the system again one to two weeks prior to harvest. Note: This step is especially important when using coco-based mediums as coco has a tendency to retain salts more than peat-based soils. Always remember to adjust your pH.

PRESCRIBED DOSAGES ARE REFLECTED IN TSP/GALLON. FEED TWO TIMES PER WEEK.

Time*	Weeks	Seedlings and Cuttings	1	2	3	4	SledgeHammer Flush: 2 tsp/gallon	5	6	7	SledgeHammer Flush: 2 tsp/gallon	8	9	10	SledgeHammer Flush: 2 tsp/gallon	11	12
Light	Hours	18	18	18	18	18			12	12		12		12		12	12
EC	Range	0.5-0.6	0.5-0.6	2.0-2.2	2.5-2.7	2.7-3.0		2.8-3.0	1.8-2.0	2.8-3.0		2.9-3.1	1.7-1.9	2.8-3.0		1.7-1.9	1.6-1.8
PPM	Range	350-420	350-420	1400-1540	1750-1890	1890-2100		1960-2100	1260-1400	1960-2100		2030-2170	1190-1330	1960-2100		1190-1330	1120-1260
Big Bloom®		6	6	6	6	3		3	3	3		3	3	3		3	3
Grow Big®				2	3	3		2		2		2		2			
Tiger Bloom®								2	2	2		2	2	2		2	2
Kangaroots™				0.5		0.5			1								
Microbe Brew™				0.5	0.5			0.5		0.5			0.5			0.5	
Open Sesame®						0.25		0.25	0.25								
Beastie Bloomz®										0.25		0.25					
ChaChing®													0.25	0.25		0.25	0.25

*For longer grow cycles, continue the week 12 feeding schedule until harvest.

